

Original article:

Study of spectrum of childhood eye diseases in a tertiary care hospital at rural Bangalore

Shrinivas Badageri

Assistant professor, Dept of ophthalmology, MVJ Medical college and Research center , Bangalore, India

Corresponding author: Dr. Shrinivas Badageri

ABSTRACT:

BACKGROUND: Eye diseases are important cause of medical consultation in children, with the spectrum varying in different localities. This study aimed to determine the spectrum of childhood eye diseases in a tertiary hospital serving rural and semi-rural communities.

METHODS: We conducted a retrospective review of all patients less than 15 years old who presented to the eye clinic of MVJ medical college between January 2011 and December 2013. Data on age at presentation, age at onset of disease, sex and diagnosis were collected and analyzed using SPSS. A *p* value less than 0.05 was considered statistically significant.

RESULTS: We evaluated the reports of 286 children, with a male:female ratio of 1:1.1. Children aged 11–15 years made up the largest group (*p*=0.013). Ocular trauma (21.7%), allergic conjunctivitis (17.8%), infections of the eye and its adnexa (15.4%) and refractive errors (14.3%) were the most common conditions. Ocular injury was more common in males (*p*=0.002) and children aged 6–10 years, and 87.1% of these cases were a closed globe injury.

CONCLUSION: The prevalent childhood eye diseases recorded here can lead to absenteeism from school and are potentially blinding. Health education aimed at the prevention of ocular trauma and prompt presentation for the management of other eye diseases should be encouraged. School health screening with periodic visual examination in the school going age group, helps in the early detection of refractive errors in children.

Keywords: Allergic conjunctivitis, Pediatric ocular disorders , Ocular trauma

INTRODUCTION

Eye diseases in children are important cause of medical consultation.¹ Children should receive prompt and proper eye care in order to avoid vision problems and eye morbidities, which could affect their learning ability, personality and adjustment in school.^{2,3} Vision is an important requirement for learning and communication.⁴ Childhood blindness is second only to cataracts in terms of “blind years.”⁵ Complaints necessitating eye care consultation can originate from

the child, parents or even teachers.⁶ The spectrum of ocular problems varies from country to country and even from region to region in the same country.² In the United States, strabismus, amblyopia and optical problems impacting visual acuity are the most common ocular problems seen among school-aged children.⁷ Refractive errors (25.7%), vernal conjunctivitis (25.3%), eye injuries (13.3%) and corneal inflammation (12.5%) were the leading causes of childhood eye morbidity reported 12 years ago in in

Ibadan, Nigeria.⁸ Meanwhile, trachoma (33.7%), refractive errors (6.3%) and non-trachomatous conjunctivitis (5.9%) were the most common childhood eye disorders reported in Ethiopia.⁹

Data on the pattern of presentation of childhood eye diseases serve as useful template for planning eye care for children in a given region. Children in rural communities may be involved in agrarian activities, which could impact their health. This study was conducted to determine the frequency and spectrum of childhood eye disorders presenting to the eye clinic of MVJ medical college Bangalore. These findings will be useful for pediatric ophthalmic care planning and could also serve as a baseline for the development of a pediatric ophthalmology subspecialty in the tertiary hospital.

METHODOLOGY

All new patients aged 15 years and younger who presented to eye clinic of MVJ medical college between January 2011 and December 2013 were retrospectively reviewed. Children who presented for a medical check-up and had no eye disorders were excluded from the study. The eye clinic of MVJ medical college serves as a referral center for the rural and semi-rural areas. It also provides primary and secondary eye care services to self-reporting patients. At the first visit, all patients had a full ophthalmic evaluation carried out by a consultant ophthalmologist. Tests were conducted to elicit a diagnosis, and management commenced as required. Subsequently, patients were seen by resident

ophthalmologists under the supervision of the consultant. Consultations to pediatricians and other specialists were made when necessary. An examination under anesthesia was carried out to confirm a diagnosis when required.

The age at presentation, sex, clinical diagnosis and age at onset of symptoms were determined from medical records. The clinical diagnosis was grouped as appropriate; diseases present at birth but not due to birth circumstances were considered congenital. Patients were grouped by age into a preschool (0–5 years), school-age (6–10 years) or older children (11–15 years) group.

Data were stored and analyzed using the Statistical Package of the Social Sciences software and are presented as simple frequencies or cross-tabulations. A chi-squared test was used to compare variables and a *p* value less than 0.05 was considered statistically significant.


RESULTS

We evaluated 286 records of children who were seen in the eye clinic during the study period, constituting 14.3% of all new patients seen. There were 138 (48.3%) males and 148 (51.7%) females, resulting in a male:female ratio of 1:1.1. Figure 1 shows the age group and sex distribution of the children seen, with a female preponderance in the 11–15 years age group. The highest frequency of consultation was recorded among older children, constituting 38.5% of the patients (*p*=0.013).

Indian Journal of Basic and Applied Medical Research


Is now with **IC Value 5.09**

Figure 1 : Age group and sex distribution of children



[Ocular injuries were the most common disorders seen (21.7%), followed by allergic conjunctivitis (17.8%), infection of the eye and its adnexa (15.4%) and refractive errors (14.3%).]

Figure 2 : shows the spectrum and frequency of eye morbidities seen. Uveitis was seen in six children, and a 15-year-old child presented with pterygium.




[The frequency and pattern of eye diseases varied across the three age groups (Table 1). The difference in presentation by age group was more prominent and statistically significant among children with refractive errors and infections. There were 29 (70.7%) cases of refractive errors among older children ($p < 0.001$), while infection of the eye adnexa was recorded more frequently (70.5%) among preschool children ($p < 0.001$)]

Table 1 : Distribution of childhood eye diseases across age groups

| Diagnosis | Age group (years) | | | |
|------------------------------|-------------------|-----------------|------------------|-----------------|
| | 0-5(%) | 6-10(%) | 11-15(%) | Total(%) |
| Ocular trauma | 16(25.8) | 26(41.9) | 20(32.3) | 62(21.7) |
| Allergic conjunctivitis | 12(24) | 12(24) | 26(52) | 50(17.5) |
| Infections of eye and adnexa | 31(70.5) | 5(11.4) | 8(18.2) | 44(15.4) |
| Refractive errors | 2(4.9) | 10(24.4) | 29(70.7) | 41(14.3) |
| Cataract | 7(36.8) | 8(42.1) | 4(21.1) | 19(6.6) |
| Corneal opacity | 4(33.3) | 4(33.3) | 4(33.3) | 12(4.2) |
| Tumors | 5(45.5) | 2(18.2) | 4(36.3) | 11(3.8) |
| Strabismus | 4(57.1) | 1(14.3) | 2(28.6) | 7(2.4) |
| Uveitis | 1(16.7) | - | 5(83.3) | 6 (2.1) |
| Glaucoma | 3(75) | - | 1(25) | 4(1.4) |
| Others | 19(63.3) | 4(13.3) | 7(23.9) | 30(10.5) |
| Total | 104(36.4) | 72(25.2) | 110(38.4) | 286(100) |

Chart Title



Among the children with ocular injury, 43 (69.4%) were boys ($p=0.002$). Ocular trauma was recorded more often among school-aged children, constituting 41.9% of all traumatic eye disorders. Eye injury was unilateral in all cases. Table 2 shows the pattern of

ocular trauma seen in the 62 children, with the majority (87.1%) being closed globe injuries. Traumatic hyphema was recorded in 5 (11.4%) of the 44 children who presented with a contusive eye injury.

Table 2 : Pattern of ocular trauma in 62 children

| Type of Injury | | Frequency (%) |
|----------------------------|------------------------------------|----------------|
| Closed globe injury | - contusion | 44(71) |
| | - superficial corneal foreign body | 9(14.5) |
| | -superior tarsal foreign body | 1(1.6) |
| Open globe injury | -penetrating | 6(9.7) |
| Others | -periorbital soft tissue injury | 1(1.6) |
| | -chemical eye injury | 1(1.6) |
| Total | | 62(100) |

Pattern of ocular trauma in 62 children

The incidence of allergic conjunctivitis was higher in older children, with 52% of the cases recorded in children aged 11–15 years ($p=0.02$). No gender differences were observed. Among the children who presented with infections, 27 (61.4%) were female ($p=0.132$). Anatomically, corneal infections accounted for 40.9% of all cases of childhood eye and adnexa infections, 33.3% of which were related

to measles All cases of corneal infection were in children aged 5 years and younger. Tumors accounted for 3.8% of the childhood eye diseases during the study period, with three cases (27.3%) of retinoblastoma, three cases (27.3%) of dermoids, two cases (18.2%) of Burkitt’s lymphoma and one case (9.1%) each of a orbital rhabdomyosarcoma, a conjunctival mass and an eye lid mass.




Figure 3 : Childhood infections of the eye and adnexa

Almost three-quarters (73.2%) of the children with refractive errors were female ($p=0.003$). Congenital eye diseases were found in 38 of the 286 (13.3%) children, of which 76.3% presented as preschool

children. Table 3 shows the pattern of congenital eye diseases seen, with congenital cataract and strabismus occurring more frequently.

Table 3 : Distribution of congenital ocular disorders among age groups

| Congenital eye disease | number | | | |
|------------------------|-----------|----------|-----------|----------------|
| | 0–5 yrs | 6–10 yrs | 11–15 yrs | Total (%) |
| Cataract | 7 | 2 | - | 9(23.7) |
| Strabismus | 4 | - | 2 | 6(15.9) |
| Glaucoma | 3 | - | - | 3(7.9) |
| Ptosis | 2 | 1 | - | 3(7.9) |
| Microphthalmia | 1 | 1 | 1 | 3(7.9) |
| Dacryostenosis | 2 | - | - | 2(5.2) |
| Presumed Leber's | 2 | - | - | 2(5.2) |
| amaurosis | | | | |
| Dermoid cyst | 1 | - | 1 | 2(5.2) |
| Eye lid eversion | 2 | - | - | 2(5.2) |
| * Others | 5 | - | 1 | 6(15.9) |
| Total | 29 | 4 | 5 | 38(100) |

[*Others are made up of one patient each with macrophthalmia, corneal opacity, persistent hyperplastic primary vitreous, oculocutaneous albinism, ocular albinism and lateral canthal skin tag.]

DISCUSSION

Children have unique problems in terms of ocular morbidities, not only due to their inability to articulate their problems, but also because of the potential to develop amblyopia in the event of visual impairment.⁴ The higher frequency of consultations in older children in this study may be due to their ability to be more articulate about their problems, while the chronicity of some of the eye disorders may also be a contributing factor. This frequency differs from a similar study reporting a higher incidence of

consultation in children younger than 5 years of age. A difference in the spectrum and incidence of childhood eye disorders may account for this.¹⁰ Ocular trauma was the most common morbidity seen in this study, which is different from other reports in which pediatric eye injuries were the third^{8,10,11} or fourth¹² most common disorder reported. In India² and other regions in Nigeria¹⁰⁻¹³, allergic conjunctivitis has been the most frequently reported childhood eye disease, while refractive error has been more frequently reported in Ibadan.⁸ The reason for

such a high frequency of pediatric eye injuries in this environment is not clear. The largely agrarian nature of the communities may predispose the children to injuries from twigs and farming activities. Further study of predisposing factors to eye injuries in children in this region is needed, given that the identification of the causes of injury helps in developing effective preventive measures.¹⁶ Eye injuries remain a major cause of unilateral visual impairment worldwide¹⁷ and a common cause of non-congenital unilateral blindness.¹⁸ Children are particularly at risk of ocular injury due to their decreased ability to detect and avoid potential hazards.^{16,19} Most childhood eye injuries are sustained during unsupervised play and domestic activities.²⁰⁻²³ The male preponderance reported here has been previously described.¹⁹⁻²³ The higher incidence of injuries reported among school children is also similar to other studies.^{16,21,24} This group represents an independent and adventurous age group, making them more vulnerable. Closed globe injury was most common, consistent with several previous studies,^{2,16,23} although a study by Kaduna found that open globe injuries were more common.²¹ The challenge of managing childhood eye injuries in this environment is enormous. Considerations range from late presentation to eye care centers to a lack of facilities, the low socioeconomic status of the children involved, the special care required during examination, postoperative management and the risk of amblyopia.^{10,16,21,23} Prevention of ocular trauma in children remains a priority in order to reduce ocular morbidity.¹⁶ This will involve the adequate education of children, parents and teachers to ensure adequate supervision at play and avoidance of risky chores. Previous reports of allergic conjunctivitis as the most common surface disorder in children^{1,10-13,25} are

corroborated by this study. Communities with largely labor and a dusty environment may be contributory.^{5,10} Moreover, rural living is a risk factor for the development of chronic allergic conjunctivitis in children.²⁶ A slight female prevalence was noted in this study, which differed from other reports that demonstrated a male prevalence.^{26,27} The higher incidence noted among older children may be related to factors affecting late presentation to an eye care facility for a chronic recurrent disease rather than the actual prevalence of the disease. Allergic conjunctivitis is a leading cause of absenteeism from school due to its discomfort, chronicity and recurrence.²⁵ Adequate management can bring symptoms under control and prevent potentially blinding complications, loss of concentration and absenteeism from school.¹⁴

Ocular infections are preventable but challenging causes of blindness, particularly in children given the number of blind years involved. Corneal infections constituted more than one-third of the infections of the eye and adnexa in this study and are particularly notorious for causing blinding corneal scars. Corneal scarring accounts for 75% of childhood blindness in Africa, with half of these cases involving a history of measles shortly before blindness.^{18,26} More recent reports have credited the Expanded Programme on Immunization for a downward trend in the incidence of measles infection and associated ocular morbidities in hosakote.^{1,10} Total eradication is desired, however, as 33% of childhood keratitis in this study was due to measles; that is, they involved a history of recent measles before the onset of eye disease. Preschool children were more likely to present with infections of the eye and adnexa in this study, and this may be related to the higher risk of measles infection in the same age group. While

efforts to establish corneal banking services are ongoing, prevention and prompt treatment of corneal infections is essential.

Refractive errors affect childhood development, given that 80% of learning in children is sight-dependent.²⁸ In the absence of regular preschool or school eye-screening for refractive errors, many children with refractive errors go unnoticed. The incidence of refractive errors in this study may have been underestimated since it was a hospital-based study, and only children with obvious complaints would be detected. Moreover, the higher incidence reported in older children could be due to better articulation and detection of visual problems by older children, suggesting a lack of detection by parents and teachers at younger ages. The statistically significant female prevalence is similar to a previous report, suggesting that young females tend to report visual problems more than males.¹⁷ School health services that regularly screen for refractive errors and refer affected children for refractive services can prevent poor performance in school and the development of amblyopia.^{3,10,14} The belief that eyeglass use in children further destroys the eye¹⁰ must be changed via health education in order to

ensure early presentation and improve the use of existing refractive services.

Nine (47.4%) childhood cataract cases in this study were congenital. Childhood cataracts are not managed as a simply smaller version of adult cataracts and pose different challenges including preoperative assessment, general anesthesia, correction of aphakia, postoperative care and follow up for posterior capsular opacification or glaucoma. Many patients are poor and cannot afford the available cataract surgical services, especially intraocular lens implantation, which is the best option for aphakic correction in most children to reduce the incidence of post-operative amblyopia.^{10,21}

In conclusion, the most common causes of childhood ocular morbidity in this study were injuries, allergic conjunctivitis, infections and refractive errors. These disorders require specialist eye care services for proper management given that they lead to absenteeism from school and are potentially blinding. Appropriate health education needs to be disseminated for the prevention of childhood eye injuries, as well as early presentation of children to eye care centers for the treatment of eye disorders.

REFERENCES

1. Nwosu SNN. Childhood eye diseases in Anambra State, Nigeria. *Nigeria Journal of Ophthalmology*. 1999;7:34–8.
2. Pratab VB, Lai HB. Pattern of pediatric ocular problem in North India. *India J Ophthalmol*. 1989;37:171–2.
3. Adegbehingbe BO, Adeoye AO, Onakpoya OH. Refractive errors in childhood. *Nigerian Journal of Surgical Sciences*. 2005;15:19–25.
4. Narayana KM, Bora A, Biswas J. Pattern of uveitis in children presenting at a tertiary eye care centre in south India. *India J Ophthalmol*. 2003;52:129–32.
5. Gilbert C. Childhood Blindness. In: Johnson G, Minassan D, Wealle R, editors. *Epidemiology of Eye Diseases*. Vol. 1998. London: Chapman and Hall; pp. 183–8.
6. King RA. Common ocular signs and symptoms in childhood. *Paed Cli North Am*. 1993;40:753–66.

7. Castenes MS. Major review: The underutilization of vision screening (for amblyopia, optical anomalies and strabismus) among preschool age children. *Binocul Vis Strabismus Q.* 2003;18:217–32.
8. Ajaiyeoba AA. Childhood eye disease in Ibadan. *Afri J Med Sci.* 1994;23:227–32.
9. Mohammed S, Abebe B. Common eye diseases in children of rural community in Goro district, Central Ethiopia. *Ethiopia J Health Dev.* 2005;19:148–52.
10. Bodunde OT, Onabolu OO. Childhood eye diseases in Sagamu. *Nigerian Journal of Ophthalmol.* 2004;12:6–9.
11. Akinsola FB. Pattern of eye diseases in Nigerian children seen at Lagos University Teaching Hospital. *Nigerian Medical Practitioner.* 1993;25:47–53.
12. Ezegwui IR, Onwasigwe EN. Pattern of eye disease in children at Abakaliki, *International Journal of Ophthalmology.* 2005;5:1128–30.
13. Abiose A. Paediatric ophthalmology problems in Lagos.. *J Trop Paediatric.* 1985;31:30–5.
14. Isawunmi MA. National Postgraduate Medical College of Nigeria; Dissertation. 2003. Ocular disorders amongst school children in Ilesa east Local Government area, Osun State, Nigeria.
15. Kuhn F, Morris R, Witherspoon D, Herman K, Jeffers JB, Trester G. A standard classification of ocular trauma. *Ophthalmology.* 1996;103:240–3.
16. Alhaski M, Almaaita J. Retrospective analysis of pediatric ocular trauma at Prince Ali Hospital. *Middle East Journal of Family Medicine.* 2007;5:42–5.
17. Niiranem M, Ratvio I. Eye injuries in children. *Br J Ophthalmol.* 1981;65:436–438.]
18. World Health Organization. Causes of Childhood blindness and current control measures. Vol. 1992. WHO; Geneva: Prevention of Childhood Blindness; pp. 21–22.
19. Olurin O. Eye injuries in Nigeria. A review of 433 cases. *Am J Ophthalmol.* 1971;72:159–66.
20. Nwosu SNN. Domestic ocular and adnexa injuries in Nigeria. *W Afr J Med.* 1995;14:137–40.
21. Kwari F, AlHassan MB, Abiose A. Pattern and outcome of paediatric ocular trauma – A 3- year review National Eye Centre Kaduna. *Nigerian Journal of Ophthalmology.* 2000;8:11–6.
22. Nkanga DG, Doln P. School vision screening in Enugu, Nigeria: Assesment of referral criteria for refractive error. *Nigerian Journal of Ophthalmology.* 1997;5:34–40.
23. Adeoye AO. Eye injuries in the young in Ile Ife, Nigeria. *Nig J Med.* 2002;11:26–28.
24. Starlman E, Elman M, Daub E, Baker S. Causes of paediatric eye injuries. A population based study. *Arch Ophthalmol.* 1990;108:376–9.
25. Wood M. Conjunctivitis: diagnosis and management. *Community Eye Health Journal.* 1999;4:7–8.
26. Bekibebe CO, Olusanya BA. Chronic Allergic Conjunctivitis: an Evaluation of Environmental Risk Factors. *Asian Journal of Ophthalmol.* 2006;8:147–50.
27. Majekodunmi S. Vernal conjunctivitis in Nigerian children. *J Peadiat Ophthalmol Strabismus.* 1978;15:176–8.
28. Yorston D. Mealses and childhood blindness. *Community Eye Health Journal.* 1991;8:2–4.
29. Sandforth-smith JH, Whittle H. Corneal ulceration following measles in Nigerian children. *Br J Ophthalmol.* 1983;63:720–4.

30. Gilbert C. Childhood Blindness: Major causes and strategies for prevention. Community Eye Health Journal. 1993;11:3-6.

Date of submission: 19 January 2014

Date of Provisional acceptance: 05 February 2014

Date of Final acceptance: 27 February 2014

Date of Publication: 04 March 2014

Source of support: Nil; Conflict of Interest: Nil