

**Original article:**

## **Assessment of Maternal Complications Associated with Caesarean Section: A Clinical Institutional Based Study**

**Suman Agarwal**

Assistant Professor, Department of Obstetrics and Gynaecology,

Teerthanker Mahaveer Medical College & Research Centre, Moradabad, Uttar Pradesh, India.

Corresponding Author: Dr.Suman Agarwal, Assistant Professor, Department of Obstetrics and Gynaecology,

Teerthanker Mahaveer Medical College & Research Centre, Moradabad, Uttar Pradesh, India.

### **Abstract**

**Background:** Caesarean section (CS) requires a longer recovery time, and operative complications such as lacerations and bleeding may occur, at rates varying from 6% for elective Caesarean to 15% for emergency Caesarean. Hence; present study was conducted to assess various maternal complications occurring in subjects undergoing CS.

**Materials & Methods:** Present study was planned to evaluate maternal complications in subjects undergoing C section. A total of 40 subjects undergoing CS were enrolled in the present study. Complete demographic details along with clinical data were obtained in all the patients. Data in relation to maternal complication was obtained and was compiled on Microsoft excel sheet. All the results were analyzed by SPSS software.

**Results:** Parity was 0 in 35 subjects while parity was 1 in 4 subjects only. Maternal complications encountered in the present study included fever, wound infection, urinary problem, peritonitis, haemorrhage, hysterectomy, anaesthetic complication, endometritis etc.

**Conclusion:** CS is associated with certain maternal complication, and therefore, it should be done only when indicated.

**Key words:** Caesarean section, Complications, Maternal.

### **INTRODUCTION**

In the present scenario, there has been an increase in the number of women who are requesting delivery by elective caesarean section without an accepted “medical indication”.<sup>1,2</sup> This trend is due in part to the general perception that caesarean delivery is much safer now than in the past and to the recognition that most studies looking at the risks of caesarean section may have been biased, as women with medical or obstetric problems were more likely to have been selected for an elective caesarean section.<sup>3-5</sup> Caesarean section (CS) requires a longer recovery time, and operative complications such as lacerations and bleeding may occur, at rates varying from 6% for elective caesarean to 15% for emergency caesarean. Having a caesarean delivery increases the risk of major bleeding in a subsequent pregnancy because of placenta previa and placental abruption. Among term babies, the risk of neonatal respiratory distress necessitating oxygen therapy is higher if delivery is by caesarean.<sup>6-8</sup>

Hence; present study was conducted to assess various maternal complications occurring in subjects undergoing CS.

### **MATERIALS & METHODS**

Present study was planned in the Department of Obstetrics and Gynaecology, Teerthanker Mahaveer Medical College & Research Centre, Moradabad, Uttar Pradesh (India) and it included evaluation of maternal complications of subjects undergoing C section. Written consent was obtained after explaining in detail the

entire research protocol. A total of 40 subjects undergoing CS were enrolled in the present study. Complete demographic details along with clinical data were obtained in all the patients.

Exclusion criteria for the present study included:

- Subjects less than 20 years of age,
- Subjects more than 35 years of age,
- Subjects with any other systemic illness

Data in relation to maternal complication was obtained and was compiled on Microsoft excel sheet. All the results were analyzed by SPSS software. One way ANOVA was used for assessment of level of significance.

## **RESULTS**

A total of 40 subjects undergoing CS were included in the present study. Mean age of the subjects of the present study was 27.5 years. 62.5 percent of the subjects of the present study were between 24 to 28 years of age. Parity was 0 in 35 subjects while parity was 1 in 4 subjects only. Maternal complications encountered in the present study included fever, wound infection, urinary problem, peritonitis, haemorrhage, hysterectomy, anaesthetic complication, endometritis etc.

## **DISCUSSION**

In the present study, the major maternal complications encountered were fever, wound infection, urinary problem, peritonitis, haemorrhage, hysterectomy, anaesthetic complication, endometritis etc. Häger RM et al determined complication rates after cesarean delivery and to identify independent risk factors for complications. The complications that were studied were intraoperative complications, blood loss, wound infection, cystitis, endometritis, hematoma, and reoperation. Independent risk factors were identified by stratification and multiple logistic regression analysis. Altogether, 21.4% of the women had  $\geq 1$  complications. The degree of cervical dilation, general anesthesia, low gestational age, and fetal macrosomia were independent risk factors. For operations that were performed at 9 to 10 cm cervical dilation, the complication rate was 32.6% versus 16.8% at 0 cm. Cesarean delivery was associated with a high complication rate. Increasing cervical dilation and, in particular, cervical dilation of 9 or 10 cm at the time of operation, general anesthesia, low gestational age, and fetal macrosomia were identified as independent risk factors.<sup>9</sup>

Loverro G et al assessed the prevalence of cesarean section (CS) related maternal complications and to evaluate post-CS complications in relationship with relative risk factors. 3010 patients who had a CS in the University Hospital of Bari during the period 1988-98 were retrospectively included into the study and 1007 women delivered vaginally at the same institution and in the same period of time, were randomly selected as the control group. In the cohort of abdominal delivery, puerperal complications were significantly more frequent compared with those following vaginal delivery ( $p < 0.05$ ). In the group of CS, obese women have higher prevalence of maternal complications, particularly hypertension and intestinal complications ( $p < 0.05$ ). Compared with vaginal delivery, CS delivery carries a higher number of postpartum complications, and the higher rate is mainly related to obesity.<sup>10</sup> Chongsuivatwong V et al compared the mortality, morbidity of emergency and elective cesarean section with vaginal delivery among Asian teaching hospitals. 12 591 vaginal deliveries, 3062 elective and 4328 emergency cesarean section were followed up to 5 days postpartum. Maternal deaths (95% CI) per 1000 births among vaginal deliveries being 0.47 was not significantly different from 0.31 of elective cesarean section and both rates were significantly lower than 2.87 per 1000 births of emergency section. The vaginal delivery group had significantly lower incidences of all major complication except significantly higher chance

of secondary operations and non-significantly different risk for endometritis. Corresponding neonatal mortality per 1000 deliveries among the three groups were 7, 2.2 and 12.4. Vaginal delivery also had higher rates of severe asphyxia and palsy than elective cesarean section. Maternal complications were increased by cesarean delivery but elective section may reduce neonatal complication.<sup>11</sup> Liu S et al assessed the maternal mortality and severe morbidity associated with low-risk planned cesarean delivery. Healthy women who underwent a primary cesarean delivery for breech presentation constituted a surrogate "planned cesarean group" considered to have undergone low-risk elective cesarean delivery, for comparison with an otherwise similar group of women who had planned to deliver vaginally. The planned cesarean group comprised 46,766 women v. 2,292,420 in the planned vaginal delivery group; overall rates of severe morbidity for the entire 14-year period were 27.3 and 9.0, respectively, per 1000 deliveries. The planned cesarean group had increased postpartum risks of cardiac arrest, wound hematoma, hysterectomy, major puerperal infection, anesthetic complications, venous thromboembolism and hemorrhage requiring hysterectomy, and stayed in hospital longer than those in the planned vaginal delivery group, but a lower risk of hemorrhage requiring blood transfusion. Absolute risk increases in severe maternal morbidity rates were low. The difference in the rate of in-hospital maternal death between the 2 groups was nonsignificant. Although the absolute difference is small, the risks of severe maternal morbidity associated with planned cesarean delivery are higher than those associated with planned vaginal delivery.<sup>12</sup>

## CONCLUSION

From the above results, it can be concluded that CS is associated with certain maternal complication, and therefore, it should be done only when indicated.

## REFERENCES


1. Matthews TG, Crowley P, Chong A, McKenna P, McGarvey C, O'Regan M. Rising caesarean section rates: A cause for concern? *BJOG* 2003;110(4):346-9.
2. Glazener CM, Herbison GP, Wilson PD, MacArthur C, Lang GD, Gee H, et al. Conservative management of persistent postnatal urinary and faecal incontinence: randomized controlled trial. *BMJ* 2001;323(7313):593-6.
3. Okada H, Kuhn C, Feillet H, et al. The 'hygiene hypothesis' for autoimmune and allergic diseases: an update. *ClinExpImmunol.* 2010;160(1):1-9.
4. McMahon MJ, Luther ER, Bowes WA, Olshan AF. Comparison of a trial of labor with an elective second cesarean section. *N Engl J Med* 1996;335(10):689-95.
5. Lumbiganon P, Laopaiboon M, Gülmezoglu M, et al. Method of delivery and pregnancy outcomes in Asia: the WHO global survey on maternal and perinatal health 2007-08. *The Lancet.* 2010;375(9713):490-9.
6. American College of Obstetricians and Gynecologists. Mode of term singleton breech delivery [ACOG Committee Opinion no 265]. *ObstetGynecol* 2001;98(6):1189-90
7. Rautava S, Walker WA. Commensal bacteria and epithelial cross talk in the developing intestine. *CurrGastroenterol Rep.* 2007;9(5):385-92.
8. Hofmeyr J, Hannah ME. Planned cesarean section for breech delivery [Cochrane review]. In: *The Cochrane Library*, Issue 3, 2003. Oxford: Update Software.

9. Häger RM1, Daltveit AK, Hofoss D, Nilsen ST, Kolaas T, Øian P, Henriksen T. Complications of cesarean deliveries: rates and risk factors. *Am J Obstet Gynecol.* 2004 Feb;190(2):428-34.
10. Loverro G1, Greco P, Vimercati A, Nicolardi V, Varcaccio-Garofalo G, Selvaggi L. Maternal complications associated with cesarean section. *J Perinat Med.* 2001;29(4):322-6.
11. Chongsuvivatwong V1, Bachtiar H et al. Maternal and fetal mortality and complications associated with cesarean section deliveries in teaching hospitals in Asia. *J ObstetGynaecol Res.* 2010 Feb;36(1):45-51.
12. Liu S1, Liston RM, Joseph KS, Heaman M, Sauve R, Kramer MS; Maternal Health Study Group of the Canadian Perinatal Surveillance System. Maternal mortality and severe morbidity associated with low-risk planned cesarean delivery versus planned vaginal delivery at term. *CMAJ.* 2007 Feb 13;176(4):455-60.

**Table 1: Demographic details of the subjects in the present study**

Parameter		Number
Number of subjects		40
Mean age (years)		27.5
Parity	0	35
	1	4
	≥2	1

**Graph 1: Age-wise distribution of subjects**



**Table 2: Distribution of subjects according to maternal complications**

<b>Maternal complication</b>	<b>Number of subjects</b>	<b>Percentage</b>
Fever	10	25
Wound infection	6	15
Urinary problem	4	10
Peritonitis	4	10
Haemorrhage	5	12.5
Hysterectomy	3	7.5
Anaesthetic complication	2	5
Endometritis	2	5
Others	2	5