

Original article

The alarming impact of obesity in inflammation by correlating with high sensitivity C-reactive protein in adults

Dr. R. Hemavathy , Dr.M.Senthilvelou, Dr.S.Chandra, Dr.M.JhansiRani, Dr.K.Soundariya

Name of the Institute/college: Department of Physiology , Sri Manakula Vinayagar Medical College and Hospital, Madagadipet, Pondicherry- 605107

Corresponding author: Dr.R.Hemavathy

Date of submission: 09 June2014; Date of Publication: 22 September 2014

Abstract:

Introduction: Obesity is primarily considered to be a disorder of energy balance, and it has recently been suggested that some forms of obesity are associated with chronic low-grade inflammation. Obesity is a chronic, multifactorial and complex disease which poses a major public health issue increasing the risk of non - communicable diseases like, Type -2-DM, Cardiovascular disease, Hypertension etc. The present study aims to emphasize the burden of obesity in inflammation by associating with elevated levels of hs CRP in adults.

Materials and methods: The study was a Hospital based cross sectional study including 30 obese individuals in the age group of 17-30 years,(15 males and 15 females)with BMI > 30 were included. Other systemic diseases were excluded. Assessment was done by Weight, Height,BMI by Quetelet index and measurement of hs CRP by Immunoturbidometer. The results were tabulated and analysed.

Results: Total of 30 obese subjects were included of which 15 were males and 15 were Females. In obese subjects hs-CRP was elevated significantly. The level of hs-CRP was proportionally elevated in relation to BMI in obese subjects.

Conclusion: Elevated hs-CRP levels in obese individuals indicates a proinflammatory state. This further concludes a risk for metabolic and cardiovascular manifestations in obese individuals.

Key words: BMI, hs CRP.

INTRODUCTION

Obesity is defined as a state of being overweight with excess body fat resulting in a significant impairment of health of a person. The most widely used method to gauge obesity is BMI which is [weight kg / height (m²)]. So BMI of 30 is the most commonly used threshold for obesity in both men and women.¹ A data from National Health and m Nutritional Examination survey (NHANES) shows that the percentage of American adult population with BMI > 30 has raised from 14.5 % to 30.5 %.¹ Approximately 55 % of adults in the

united States are overweight, and nearly 22 % of adults are obese.²

Obesity is a long term disease having many serious consequences for health and it is a leading cause of preventable death in united states.³ Obesity has become so common in developed and developing nations that presently it replaces undernutrition and infectious diseases as the most significant contributor to ill health.⁴ Few decades before, obesity was considered as an indication of wealth and health or maximally as a cosmetic problem. But, now it has become an epidemic that threatens

global well being even in growing children and adolescents.⁵ Genes that predispose to obesity in humans and animals have already been identified - the faulty gene -FTO, indicating the importance of genetic factor in the development of this disorder.⁶ Recent research has shown that the number as well as the size of adipocytes in obese people is more than people with normal body weight. Weight loss in an obese adult can reduce the size but not the number of adipocytes.⁷ Hence these adults do retain the tendency to put on weight faster than normal individuals. Therefore, prevention as well as early intervention is crucial to fight obesity.

CRP is a trace protein normally present in the circulation of healthy subjects, with a medium concentration of 1 mg/ L increasing 1000 fold in response to infection, ischaemia, trauma, burns and inflammatory conditions and decrease just as rapidly with the resolution of the pathology.⁸ Hence, it is an acute phase protein which gets elevated during episodes of acute-inflammation or infection.⁹ It is synthesized by the liver in response to factor released by macrophages and fat cells.¹⁰ The physiologic role of CRP is to bind to phosphocholine expressed on the surface of dead or dying cell (and some types of bacteria) in order to activate the complement system via the C1q complex as an immunologic defense mechanism.¹¹ The present torrent of studies of C – reactive protein in Cardio vascular disease and associated conditions is facilitated by the ready commercial availability of automated CRP assays and of CRP itself as a research reagent. In the mid 1990's, immunoassays for CRP, with greater sensitivity than those previously in routine use, revealed that increased C- reactive protein values, within the range previously considered normal, strongly predicts future coronary events.¹² The high sensitivity CRP is considered as an novel biomarker, since it measures even low levels of

CRP in human serum or plasma.¹³ Obesity, mainly visceral, often associates with co-morbid conditions, like insulin resistance, glucose and lipid abnormalities, and hypertension, each one an independent cardiovascular risk factor per se.¹⁴ Obesity is currently regarded as a proinflammatory state. Excess adiposity often correlates with abnormal production of several mediators which often associate with cardiovascular events. The adipokine imbalance characterizing obesity, including low levels of adiponectin, high levels of leptin, inflammatory mediators (IL-6 and TNF- α) and antifibrinolytic factors (PAI-1) may induce oxidative stress and endothelial dysfunction- initial steps of atherogenesis.¹⁵

Numerous studies also support an independent association between circulating levels of CRP, an inflammatory marker potently induced by IL-6 and TNF- α (in excess in the obesity state), and cardiovascular events. Many studies have suggested that higher BMI is associated with higher CRP concentrations indicating a state of low – grade inflammation. Elevated CRP levels were present in 27.6% women and 6.7 % men of US adult population in obese persons with BMI > 30.¹⁶

As many studies have been conducted regarding inflammatory markers related with Cardio vascular diseases, Diabetes mellitus, Hypertension, Metabolic syndrome, Cancer, Sleep – Breathing disorders, Chronic obstructive pulmonary diseases, Bronchial asthma, Post menopausal women, PCOD's, Statin therapy etc., Only a few studies have been done based on the impact of hs CRP (High Sensitivity C – reactive protein) prevalence in obesity.

Due to the paucity of information for correlation between BMI and hs CRP in Indian settings, the present study was undertaken. Elevated hs-CRP levels in young obese individuals is an indication of an existing pro-inflammatory state in them, which

might lead to complications in future. Hence this study may create awareness among young obese individuals regarding the risk factors of obesity and the importance of diet, life style modifications and physical activities and thereby help them to lead a healthy life. Thus the present study aimed to correlate the concentration of hs-CRP in obese individuals.

Aim: : The present study aims to emphasize the burden of obesity in inflammation by associating with elevated levels of hs CRP in adults.

MATERIALS AND METHODS

Study was conducted at Sri Manakula Vinayagar Medical College and Hospital, Madagadipet, Puducherry between the period October 2012 to August 2013 after getting approved from the Institutional Research Council and Ethical committee. The subjects included into the study were selected after obtaining informed consent on the basis of fulfillment of inclusion criteria whilst excluding exclusion criteria. The study was a cross sectional study. 30 obese individuals such as Staff, Students, Persons attending medicine OPD (15 males and 15 females) in the age group of 17-30 years with BMI > 30 constituted the study population. other systemic diseases like diabetes, hypertension, hyperlipidemia were excluded.

PROCEDURES:

BODY MASS INDEX :

For the assessment of obesity, height and weight measurements were taken on each subject, and using Quetelet index, the value of BMI was calculated for each subject.

$$\text{BMI} = \text{weight (kg)} / \text{height (m)}^2$$

Classification of BMI:

Underweight : 18.5

Normal weight: 18.5-24.9

Overweight : 25-29.9

Obesity :

Class 1 : 30-34.9

Class 2 : 35-39.9

Class 3: ≥ 40

HEIGHT : It was measured against a vertical board with an attached metric rule and a horizontal headboard was brought in contact with the upper most point on the head. It was recorded barefoot, with person standing on a flat surface and weight distributed evenly on both feet and heels together and the head positioned so that the line of vision is perpendicular to the body. The arms were hanging freely by the sides and the head, back, buttocks and heels were in contact with vertical board. The individual was asked to inhale deeply and maintained a full erect position. The head was held erect with the external auditory apparatus and the lower border of the orbit in one horizontal plane. Top most point on the head with sufficient pressure to compress the hair was taken as height to the nearest of 0.1 cm.

WEIGHT : It was measured with subjects in light clothes and without shoes in erect posture using standard apparatus. weight was measured to the nearest 0.1 kg on a weighing machine (Electronic Weighing machine).

HIGH SENSITIVITY C-REACTIVE PROTEIN:¹⁷
IMMUNOTURBIDOMETER (BEACON) :
Version 3.81.¹⁷

Principle:¹⁷

The analyzer is designed on the basis of Lambert-Beer Law. The hs- CRP Turbilatex test is a quantitative turbidimetric test for the measurement of low levels of CRP in human serum or plasma. Latex particles coated with specific anti-human CRP were agglutinated when mixed with samples containing CRP. The agglutination caused an absorbance change dependent upon the CRP contents of the patient sample that can be

quantified by comparison from a calibrator of known CRP concentration.

REFERENCE VALUES :

Below 3 mg/ L is considered as normal.

PROCEDURE:

2 ml of blood was taken in a test tube and left undisturbed for half an hour for complete clot formation. The sample was then centrifuged at 3000 rpm to separate the serum from the clot. After centrifugation, the serum was stored at 20° C in Ependorf tubes till the analysis for hs-CRP was conducted. Serum hs-CRP levels were measured by turbidimetry method, using commercial kit (Beacon Diagnostics Pvt Ltd, India). hsCRP kit : Code no -T05, Pack size - 32 ml.

Statistical analysis: At 95 % CI, 80% Power, minimum sample needed is 30.¹⁸The values of mean ± SD for the obese subjects to evaluate hs CRP levels are 1.710 ± 0.3434 and 2.450 ± 0.5130 respectively. The data collected was entered in Microsoft Excel and analyzed using SPSS software package Version 19. Pearson correlation was used to analyse the hs CRP thresholds with BMI of obese individuals.

RESULTS

Total of 30 obese individuals were selected for the study of which 15 were males and remaining 15 were females categorised by calculating their BMI. The results were tabulated and analysed.

Table 1: Age and gender distribution in obese subjects.

Age (years)	Females n (%)	Males n (%)	Total n (%)
20-24	6 (40%)	7 (47%)	13 (43.3%)
25-27	3 (20%)	4 (27%)	7 (23.3%)
28-30	6 (40%)	4 (27%)	10 (33.3%)

Among these individuals 43.3% were between the age 20-24 yrs, 23.3% were between 25-27 yrs of age and 33.3% were between the age 28-30 yrs.

Table 2: BMI and gender distribution in study subjects.

BMI	Females n (%)	Males n (%)
30-34.9	9 (60%)	9 (60%)
35-39	5 (33.3%)	6 (40%)
≥ 40	1 (7%)	0 (0%)


Among the 30 obese individuals (15 Females and 15 Males) whose BMI was categorized as Class I 30 to 34.9 Kg/ m², Class II 35 to 39.9 Kg/ m², Class III ≥ 40 Kg/ m² respectively. Among them 60% of both males and females had BMI of Class I, 40% males and 33.3% females had BMI of Class II and 7% of females and none of the males had BMI of Class III which showed no significant association between BMI and gender distribution in obese individuals.

Table 3: Association between hs CRP and gender in subjects.

Reported hs CRP (mg/L)	Females n(%)	Males n (%)
< 3	3 (20%)	6 (40%)
≤ 3	12 (80%)	9 (60%)
Total	15 (100%)	15 (100%)

On the basis of gender, all the individuals included in this study were divided in two groups and were screened for the hs CRP levels. Total of 15 females and 15 males were screened and found that 80% females showed elevated hs CRP, as compared to 60% of males. Gender specificity was assessed for their association with hsCRP levels and no statistical significant association was reported (chi square 1.67 with a p value of > 0.5 NS

Figure 1: Scatter plot to show relation between BMI and hs CRP in study subjects.



A total of 30 individuals were screened for the strength of association of obesity with high sensitivity CRP. Based on the BMI calculation they were categorized as obese subjects which showed that, there was a significant relationship between BMI and levels of hsCRP, Pearson correlation coefficient $r = 0.59$, $P = 0.01$. Obesity was associated with high values of hsCRP.

All the individuals included in this study were also interviewed for the risk factors i.e. hypertension, hyperlipidemia, diabetes mellitus, drug intake,

alcohol and smoking and found that these conditions did not illustrate any relation with obesity and levels of hs CRP.

DISCUSSION

Obesity is a long term disease having many serious consequences on the health of a person.² It is a state of excess adipose tissue mass and the most widely used method to calculate obesity is BMI.¹ Recently it has also been described as a pro inflammatory state which is involved in the pathophysiology of many diseases.

The main aim of the present study was to explore the association between the body mass index and hsCRP level among the obese individuals.

Total of 30 obese individuals were selected for the study of which 15 were males and remaining 15 were females categorised by calculating their BMI. Among these individuals 43.3% were between the age 20-24 yrs, 23.3% were between 25-27 yrs of age and 33.3% were between the age 28-30 yrs. Among the 30 obese (15 males and 15 females) individuals who were subjected to relative BMI by using Quetelet index, no statistical significant difference in BMI was observed among males and females. The similar observation was made by other studies determined by Deepa M¹⁹ and her colleagues and Desigamani Kanniyappan²⁰ and his colleagues.

The high sensitivity C- reactive protein is considered as the marker of inflammation inside the body. In this cross sectional study all obese individuals were analysed for the level of hsCRP and BMI and found that hsCRP was significantly elevated among the obese individuals. In this study

the gender wise distribution of hs CRP was calculated among the controls and obese population and found that hs CRP was not significantly elevated in any specific gender in accordance with obesity. Our results, together with the evidence of previous studies by Visser M et al²¹, Hussain S D²², Preethi B L²³, Chieh Lin C et al²⁴, Shilpa B A et al²⁵, Rogowski O et al²⁶ indicates the overweightness and obesity as a risk factor for one's own health.

The obesity causes the structural and functional changes in various body sites i.e. skeletal muscle, liver, cardiovascular, integument, pulmonary, cerebral and coronary arteries. In recent reports obesity has linked with development of fatty liver diseases.²⁷ The person with obesity is associated with elevated CRP which showed frequent episodes of infections. In obesity there is increased secretion of IL - 6 which activates the hepatic pathways to release the large amount of acute-phase proteins CRP in circulation.²⁸ The older experimental studies and cross-sectional studies have showed that CRP along with IL - 6 are contributing in the development of hyperglycemic, insulin resistance and Type 2 - DM.²⁹

CRP a liver derived pentraxin, also correlates well with other risk factors and increased cardiovascular risk in the absence of acute inflammation. In recent years, it has emerged as one of the most promising biomarker for future cardiovascular events and peripheral vascular diseases in the obese individuals.³⁰ CRP which might promote the formation of intimal neovessels in vulnerable atherosclerotic plaques results in increasing the likelihood of rupture.³¹ Many research studies has tested the utility of CRP level as a marker for the initiation and monitoring of treatment with statins for obese individuals.³² Obesity activates the pathways for the production of abnormal adipokine and cytokines which can be

considered as biological markers of inflammation.³³

The link between obesity and inflammation is based on two basic theories. According to first theory, obesity - induced inflammation has been considered as a protective mechanism, which stops the body from losing activity or fitness by storing the fat in tissues and organs by anabolic process. Second theory suggest that inflammation as an catabolic process which break down the organs and tissue to control the body weight within the normal limits.³⁴

The accumulation of macrophage and lymphocyte in adipose tissue might contribute to the pathogenesis of obesity associated disorders but in recent research studies it has been found that adipose tissue acts as a secretory organ, which synthesizes various hormones, peptides and cytokines, which are involved in food intake regulation, inflammation, coagulation and blood pressure control, glucose and lipid metabolism.³⁵ All the individuals included in this study were also interviewed for the risk factors i.e. hypertension, hyperlipidemia, diabetes mellitus, drug intake, alcohol and smoking and found that these conditions did not illustrate any relation with obesity and levels of hs CRP.³⁶

In obese individuals the non pharmacological methods such as weight loss, food habits and regular exercise may help in controlling the CRP to attain a normal level, which may help to reduce the risk of development of various diseases.

CONCLUSION

The present study highlights the problem of obesity in the society which may impact on the overall performance of obese people, which implies on health as well as economical policies of the society. Elevated hs CRP levels in young obese individuals heightens the inflammatory link, which might lead to drastic complications in the future inducing the progression of non - communicable diseases

resulting in high mortality among the young population. Higher the BMI, the levels of hs CRP increases, suggesting that obesity represents a major risk factor and constitutes a serious threat to the current and future health of all population on earth, in which inflammation acts as a major driver in the pathogenesis of obesity.

In obese individuals the non pharmacological methods such as weight loss, food habits and regular exercise may help in controlling the CRP to attain a normal level, which may help to reduce the risk of development of various diseases.

The present study advocated strongly to encourage people for healthy dietary patterns, regular assessment of BMI and biochemical parameters.

REFERENCES

1. Fauci A, Braunwald E, Kasper L, Larry L. Harrison's principles of internal medicine. 18th edition. USA: McGraw Hill; 2012; Vol (1), Chapter 77 and 78, Biology of Obesity, Evaluation and Management of Obesity. P 622-636.
2. Guyton AC, Hall JE. Text Book of Medical Physiology. 12th ed. Noida: Elsevier; 2012; Chapter 71, Metabolism and Temperature Regulation. P 850-851.
3. Must A, Spadano J, Coakley EH, Field AE, Colditz G, Dietz WH. The Disease Burden Associated with Overweight and Obesity. JAMA 1999; 282:1523-1527.
4. Shah NR, Braverman ER. Measuring adiposity in patients: The utility of Body Mass Index (BMI), Percent body fat, and Leptin. PLOS ONE 2012; 7(4):e 33308.
5. Pasco JA, Nicholson GC, Brennan SL, Kotowicz. Prevalence of Obesity and the relationship between the Body Mass Index and Body Fat: Cross-Sectional, Population-Based Data. PLOS ONE 2012; 7(1):e 29580.
6. Wu J, Xu J, Zhang Z, Ren J, Li Y, Wang J et al. Association of FTO polymorphisms with obesity and metabolic parameters in Han Chinese adolescents. PLOS ONE 2014; 9(6):e 98984
7. Kershaw EE, Flier JS. Adipose tissue as an endocrine organ. The journal of clinical Endocrinology and Metabolism 2004; 89(6):2548-2556.
8. Pepys MB, Baltz ML. Acute phase proteins with special reference to C-reactive protein and related proteins (pentraxins) and serum amyloid A protein. Adv Immunol 1983; 34:141-212.
9. Lau DC, Dhillon B, Yan H, Szmitko PE, Verma S. Adipokines: molecular links between obesity and atherosclerosis. AMJ Physiol Heart Circ Physiol 2004; 288(5):2031-2041.
10. Van DM, De Maat MP, Hak AE, Kiliaan AJ, Delsol AI, Vander Kuip et al. CRP predicts progression of atherosclerosis measured at various sites in the arterial tree. Stroke 2002; 33:2750-2755.
11. Thompson D, Pepys MB, Wood SP. The physiological structure of human C-reactive protein and its complex with phosphocholine. Structure 1999; 7(2):169-177.
12. Pepys MB, Hirschfield GM. C -reactive protein: a critical update. J Clin Invest 2003; 111(12):1805-1812.
13. Napoli MD, Shwaninger M, Cappelli R, Ceccarelli E, Donati C. Evaluation of C-reactive protein measurement for assessing the risk and prognosis in ischemic stroke: A statement for health care professionals from the CRP pooling project members. Stroke 2005; 36:1316-1329.
14. Gregor MF, Hotamisligil GS. Inflammatory mechanisms in obesity. Annal Review of Immunology 2011; 29:415-445.

15. Van Gaal LF, Mertens IL, De Block CE. Mechanisms linking obesity with cardiovascular disease. *Nature* 2006; 444(7121):875-880.
16. Alley DE, Seeman TE, Kikim J, Karlamangla A, Hup et al. Socioeconomic status and C-reactive protein levels in the US population: NHANES IV. *Brain Behav Immun* 2006; 20:498-504. . Chetana 17. Vaishnavi. *Immunology and Infectious Diseases* 1996; 6:139-144.
18. Rajendran K, Devarajan N, Ganesan M, Ragnathan M. Obesity, inflammation and acute myocardial infarction - expression of leptin, IL-6 and high sensitivity - CRP in Chennai based population. *Thrombosis Journal* 2012; 10:13.
19. Deepa M, Farooq S, Deepa R, Manjula D, Mohan V. Prevalence and significance of generalized and central body obesity in an urban Asian Indian population in Chennai, India (CURES: 47). *European Journal of Clinical Nutrition* 2009; 63:259-267.
20. Kanniyappan D, Kalidhas P, Mary Aruna R. Age, gender related prevalence of cardiovascular risk factors in overweight and obese South Indian Adults. *International Journal of Biological and Medical Research* 2011; 2(2):513-522. .
21. Visser M, Bouter LM, McQuillan GM, Wener MH, Harris TB. Elevated C-reactive protein levels in Overweight and Obese Adults. *JAMA* 1999; 282(22):2131-2135.
22. Hussain SD. The correlation between serum high sensitivity C-reactive protein and leptin in reproductive age Overweight/ Obese women in Erbil city. *Zanco J Med Sci* 2012; 16(3):167-175.
23. Preethi BL, Jaisri G. Risk stratification of Body Mass Index. *National Journal of Medical Research* 2014 4(1):1-6.
24. Chieh Lin C, Kardias SLR, Ing Li C, Shong Liu C, May Lai M, Yuan Lin W et al. The relationship of high sensitivity C-reactive protein to percent body fat mass, body mass index, waist-to-hip ratio and waist circumference in a Taiwanese population. *BMC Public Health* 2010; 10:579.
25. Shilpa BA, Jayashree SB, Amruta M, Mangesh T, Balaji NA. High sensitivity C-reactive protein : An independent proinflammatory cardiac marker in Healthy Overweight and Obese individuals. *Journal of Research in Obesity* 2014; 17:1-8.
26. Rogowski O, Shapira I, Toker S, Melamad S, Shiron A, Berliner S et al. Obesity-related correlation between C-reactive protein and the calculated 10-Y Framingham Coronary Heart Disease Risk Score. *International Journal of Obesity* 2005. 29:772-777.
27. Farrell GC, Larter CZ. Nonalcoholic fatty liver disease : from steatosis to cirrhosis. *Hepatology* 2006; 43(2 supplement 1):S99-S112.
28. Yudkin JS, Stehouwer CD, Emeis JJ, Coppack SW. C-reactive protein in healthy subjects : associations with obesity, insulin resistance and endothelial dysfunction : a potential role for cytokines originating from adipose tissue? *Arteriosclerosis, Thrombosis and Vascular Biology* 1999; 19(4):972-978.
29. Sandler S, Bendtzen K, Eizirik DL, Welsh M. Interleukin-6 affects insulin secretion and glucose metabolism of rat pancreatic islets in vitro. *Endocrinology* 1990; 126(2):1288-1294.
30. Ridker PM, Cushman M, Stampfer MJ, Tracy RP, Hennekens CH. Inflammation, aspirin and the risk of cardiovascular disease in apparently healthy men. *New England Journal of Medicine* 1997; 336(14):973-979.

31. Ridker PM, Stampfer MJ, Rifai N. Novel risk factors for systemic atherosclerosis : a comparison of C-reactive protein, fibrinogen, homocysteine, lipoprotein (a) and standard cholesterol screening as predictors of peripheral arterial disease. *Journal of the American Medical Association* 2001; 285(19):2481-2485.
32. Ridker PM, Danielson E, Fonseca FAH et al. Rosuvastatin to prevent vascular events in man and women with elevated C-reactive protein. *New England Journal of Medicine* 2008; 359(21):2195-2207.
33. Hotamisligil GS, Shargil NS, Spiegelman BM. Adipose expression of tumor necrosis factor α : direct role in obesity-linked insulin resistance. *Science* 1993; 259(5091):87-91.
34. Greenberg AS, Obin MS. Obesity and the role of adipose tissue in inflammation and metabolism. *The American Journal of Clinical Nutrition* 2006; 83(2):461S-465S.
35. Hajer GR, Van Haeften TW, Visseren FLJ. Adipose tissue dysfunction in obesity, diabetes and vascular diseases. *European Heart Journal* 2008; 29(24):2959-2971.
36. McLaughlin T, Abbasi F, Lamendola C, Liang L, Reaven G, Schaaf P et al. Differentiation between obesity and insulin resistance in the association with C-reactive protein. *Circulation* 2002; 106:2908-2912.