

Original article

Physiological effect of substance abuse on the Pulmonary

Functions in rural Uttar Pradesh

Rubeena Bano¹, Sanjay Khanna², Nadeem Ahmad³, Siddiqi MN⁴

¹Assoc. Professor, Dept. of Physiology, ²Assoc. Professor, Dept. of Pharmacology, ³Professor, Dept. of Community Medicine, ⁴Assoc. Professor, Dept. of Forensic Medicine, Integral Institute of Medical Sciences & Research (IIMS&R), Lucknow, U.P.

Correspondence to: Dr. Rubeena Bano

Abstract:

Background: In India substance abuse is a common habit prevalent in both urban and rural areas. Cigarette and bidi smoking, tobacco chewing, has extensive effects on respiratory function and is clearly implicated in the etiology of a number of respiratory diseases.

Objectives: 1. To study and compare the pulmonary function tests among smokers and non-smokers in a rural area. 2. To study the role of possible associated factors and relation of type, quantity and duration of common substances abused on the pulmonary function tests.

Methods: A cross sectional study was carried out in Hind Hospital, Lucknow to assess the pulmonary function tests on computerized spirometer in 200 male subjects, having habit of substance abuse, in field practice area of HIMS, Lucknow. **Statistical analysis:** SPSS Statistical Software.

Results: Almost all the pulmonary function parameters were significantly reduced in smokers as compared to other substances of abuse liability and obstructive pulmonary impairment was commonest.

Conclusion: By spirometry a spectrum of pulmonary disorders may be detected at an early stage and subsequent morbidity can be minimized.

Key words: Substance abuse, Spirometry, Pulmonary functions, Rural area

Introduction:

Substance use refers to the use of any psychoactive substances or drugs, which include licit and illicit drugs, other than which are medically indicated.[1,2] Psychoactive substance use poses a threat to the health and social and economic fabric of families, communities and nations.[3] Drug dependence is a growing problem, consequences of drug dependence cost the community heavily and they form a major health problem.[4] This habit not only affects health, education and occupational career, but it

also incurs a huge financial and social burden on the society.[5]

The epidemic of substance abuse in young generation has assumed alarming dimensions in India. Changing cultural values, increasing economic stress and dwindling supportive bonds are leading to initiation into substance use. According to the World Health Organization (WHO) substance abuse is persistent or sporadic drug use inconsistent with or unrelated to acceptable medical practice.[6] About 190 million people all over the world consume one

drug or the other.[7] This Drug addiction causes immense human distress and the illegal production and distribution of drugs have spawned crime and violence worldwide. Cannabis, heroin, and Indian-produced pharmaceutical drugs are the most frequently abused drugs in India. Cannabis products, often called charas, bhang, or ganja, are abused throughout the country.[8] In India smoking is a common habit prevalent in both urban and rural areas irrespective of mode of smoking i.e. cigarettes, bidis, pipes, cigar, hookah etc. The Adolescent drug abuse is one of the major areas of concern in adolescent and young people's behavior. It is estimated that, in India, by the time most boys reach the ninth grade, about 50 percent of them have tried at least one of the substance of abuse nature.[9] In last three decades, many epidemiological surveys have been carried out in India to assess the prevalence of substance abuse.

Materials and Methods:

The present cross sectional study was conducted in Hind Hospital of Lucknow district, Uttar Pradesh. The study population included 200 male subjects having habit of substance abuse, of one type or the other, aged between 30-60 years. Individuals with history of smoking cigarettes / bidis daily for at least one year were considered

as smokers. The materials used in the study were a computerized RMS Med-spirometer, weighing machine, measuring tape and Blood Pressure set. The observations of the study were analyzed by statistical methods like percentages, chi square test and t-test of significance.

Results:

In the present study it was observed that there was no significant difference in the mean physical parameters like age, height, weight, body mass index and body surface area by calculating mean and standard deviation in different substance abusers (**Table 1**).

Most of the smokers smoked only bidi (52.0%) followed by only cigarettes (9.0%). Tobacco chewers (Gutkha, Panmasala, Khaini, Pan, Mishri etc.) were 24.0%, 12.0% were alcoholics and only 3.0% were involved in hard drug abuse (Cannabis, i.v.drug use, opium, ganja, brown sugar etc.) (**Table 2**).

All Pulmonary function parameters like FVC, FEV₁, FEV₁/FVC, PEFR, FEF_{25-75%} and MVV showed statistically highly significant association between smokers and non-smokers by applying unpaired t-test of significance ($p < 0.001$). The obstructive lung changes were most common and were observed predominantly in smokers (26.27%) as compared to other substance abusers (**Table 3**).

Indian Journal of Basic and Applied Medical Research

Is now with

IC Value 5.09

Table 1: Physical variables of substance abusers

Variables	Smoking Mean ± 2 S.D.	Tobacco chewing Mean ± 2 S.D.	Alcohol Mean ± 2 S.D.	Others (Cannabis, charas, bhang, or ganja) Mean ± 2 S.D.
Age (years)	48.26± 10.09	48.10 ± 10.54	45.15 ± 12.14	46.18 ± 13.57
Height (m)	1.66 ±0.11	1.67 ± 0.12	1.62 ± 0.32	1.61 ± 0.62
Weight (Kg)	65.4 ± 8.8	64.4 ± 11.5	61.4 ± 13.7	60.4 ± 2.5
Body Mass Index	23.52 ± 3.20	23.80 ± 3.37	22.51 ± 2.68	24.86 ± 2.30
Body surface area	1.71 ± 0.06	1.74 ± 0.14	1.54 ± 0.23	1.68 ± 0.48

S.D. = Standard Deviation

Table 2: Distribution of substances abused

Category of substances abused	No.	%
Bidi smoking	104	52.0
Cigarette smoking	18	9.0
Tobacco chewing (Gutkha/Pan masala/Pan/Khaini/Mishri)	48	24.0
Alcohol	24	12.0
Others (Cannabis products, charas, bhang, or ganja, etc.)	6	3.0
Total	200	100.0

Table 3: Interpretation of pulmonary function tests (PFT) in substance abusers

PFT Results	Smokers No. (%)	Tobacco chewers No. (%)	Alcoholics No. (%)	Others No. (%)
Obstructive	32 (26.27)	11 (22.91)	2 (8.33)	1 (16.66)
Restrictive	8 (6.58)	4 (8.33)	2 (8.33)	1 (16.66)
Mixed	18 (14.76)	2 (4.16)	1 (4.16)	0 (0.0)
Normal	64 (52.45)	31 (64.58)	19 (79.16)	4 (66.66)
Total	122 (100.0)	48 (100.0)	24 (100.0)	6 (100.0)

Chi square value = 20.84, p<0.001, highly significant.

Discussion:

In the present study it was observed that there was no significant difference in the mean physical parameters like age, height, weight, body mass index and body surface area of substance abusers. None of individuals smoked tobacco in any form other than bidis or cigarettes. Most smokers were bidi smokers. In India, tobacco is consumed mainly in the form of bidis, followed by smokeless tobacco and cigarettes. Bidi smoke may be more injurious because bidi contains unrefined form of tobacco as compared to cigarettes. Cigarette smoking has extensive effects on respiratory function and is clearly implicated in the etiology of a number of respiratory diseases, particularly chronic bronchitis, emphysema, and bronchial carcinoma. All Pulmonary function parameters showed statistically highly significant association between smokers and non-smokers by applying unpaired t-test of significance ($p < 0.001$). In Uttar Pradesh, Dube and Handa [10] reported that 22.8 per 1000 were dependent on alcohol and drugs while Thacore [11] from Lucknow gave a figure of 18.55 per 1000. Various epidemiological surveys also revealed that 20-40% of subjects above 15 years are current users of alcohol and 10% of them are

regular or excessive users.[12,13] In a rural population of Uttar Pradesh alcohol was found to be the commonest substance abused (82.5%) followed by cannabis (16.1%). Varma et al [14] found that rates of current use of alcohol in Punjab were 45.9% in Jalandhar and 27.7% in Chandigarh whereas it was 28.1% in rural areas of Punjab.[15] Shukla [16] reported that 38.3% of the rural population in Uttar Pradesh was habitual substance users. In a study conducted in rural community in Bihar prevalence of alcohol/drug use was found to be 28.8% of the study population.[17]

Conclusion:

The pulmonary function tests were assessed on a computerized spirometer in 200 male substance abusers. The present study reveals the effect of type, duration and pattern of substance abuse in rural community of India. Bidi smoking was most common as the study setting was in rural India. Almost all the pulmonary function parameters were significantly reduced in smokers as compared to non smoker substance abusers and obstructive pulmonary impairment was commonest in all the substance abusers. Efforts should be made to educate the community about the harmful effects of substance abuse.

References:

1. World Health Organization. Guide to drug use epidemiology. Technical Report Series, 2012.
2. Smart RG, Hughes PH, Johnston LD, Anumonye A, Khant U, Medina Mora ME, et al. A methodology for student drug-use surveys. Geneva: World Health Organization. 1980.
3. Swadi H. Drug abuse in children and adolescents: An update. Arch Dis Child. 1992;67:1245-46.
4. Ramachandran V. The prevention of alcohol related problems. Indian J Psychiatry. 1991;33:3-10.
5. Ningombam S, Hutin Y, Murhekar MV. Prevalence and pattern of substance use among the higher secondary school students of Imphal, Manipur, India. Natl Med J India. 2011;24:11-15.

6. WHO. Techn. Res. Ser. No.886: 1999. p. 48.
7. Miller, WR. Sanchez, VC. Motivating young adults for treatment and lifestyle change. 1993. pp. 55-82.
8. Srivastava A. Pal HR. Dwivedi SN et al. National household survey of drug abuse in India. Report submitted to the Indian Ministry of Social Justice and Empowerment and the United Nations Office for Drugs and Crime. 2003.
9. Ramachandran V. The prevention of alcohol related problems. Indian J Psychiatry 1991; 33:3-10.
10. Dube KC, Handa SK. Drug use in health and mental illness in an Indian population. Br J Psychiatry 1971;118:345-46.
11. Thacore VR. Drug abuse in India with special reference to Lucknow. Indian J Psychiatry 1972;14:257-61.
12. Dube KC, Kumar A, Kumar N, Gupta SP. Prevalence and pattern of drug use amongst college students. Acta Psychiat Scand 1978; 57:336-46.
13. Lal B, Singh G. Drug abuse in Punjab. Br J Addict 1979; 74:441.
14. Varma VK, Singh A, Singh S, Malhotra AK. Extent and pattern of alcohol use in North India. Indian J Psychiatry 1980; 22:331-37.
15. Lal B, Singh G. Alcohol consumption in Punjab. Indian J Psychiatry 1978; 20:212-16.
16. Shukla BR. Drinks and drugs in a north Indian village-an anthropological study. Ethnographic and Folk culture Society: Lucknow, India; 1979.
17. Jena R, Shukla TR, Hemraj P. Drug abuse in a rural community in Bihar: Some psychosocial correlates. Indian J Psychiatry 1996;38:43-6.