Review article:

Nutritional benefits and pharmacological effects of ginger: an overview

Dr. (Mrs.) Anita Singh

Assistant Professor and head, Department of Home Science Kr. R.C.M. P.G College Mainpuri, U.P- 205001

Abstract

Ginger the rhizome of the Zingiberofficinale is used worldwide as a cooking spice, condiment and herbal remedy. It is primarily used as an anti-inflammatory, a pain remedy, a warming remedy and a cholesterol-lowering herb. Randomized controlled trials support its use in preventing nausea. Case studies suggest usefulness in treating migraines and inflammatory arthritis, but no randomized trials have been reported. Given its long history of use as a food, ginger is presumed safe for supplemental use. Fresh ginger contains 80.9% moisture, 2.3% protein, 0.9% fat, 1.2% minerals, 2.4% fibre and 12.3% carbohydrates. The active components of ginger are reported to stimulate digestion, absorption, relieve constipation and flatulence by increasing muscular activity in the digestive.

Keywords: Anti-inflammatory, Ginger, cholesterol-lowering, neuro-protective.

Introduction

Ginger has been used by traditional Chinese and Indian medicine for over 25 centuries. [1] Ginger was brought to Mexico by the Spaniards and later introduced to Jamaica, the latter currently being one of the world's foremost producers of this species. [2] In recent times, ginger has been introduced into various tropical countries where diverse chemotypes have been developed. [3] Ginger (Zingiberofficinale Roscoe) is a member of the Zingiberaceae family of plants. The English term 'ginger' originated from Sanskrit word 'Sringavera' which means horn-like. The underground stem (rhizome) is used for preparation of ginger. This rhizome can be processed into a powder, syrup, volatile oil, and oleoresin. The rhizome contains fats, carbohydrates, protein, fiber, water, and volatile oil. It has been a part of healing strategies in Asia, India, Europe, and the Middle East for centuries for treatment of such disorders as arthritis, stomach upset, asthma, diabetes, and menstrual irregularities, to name a few. [4] According to a 2010 study published in the "Journal of Microbiology and Antimicrobials", ginger proved to have higher antimicrobial power than conventional antibiotics against two strains of staph infections. Ginger is thought to have anti-inflammatory properties, sometimes used to treat arthritis. Ginger has been used for its herbal properties, which are especially helpful in easing stomach and motion sickness. This herb has been effective in controlling nausea and vomiting. It is hypothesized to work by changing serotonin receptors in the digestive tract. Ginger appears to work like ibuprofen for menstrual pain, according to one of the study. [5]

The main aim to write this review is to give insight on *Zingiberofficinale* about its valuable nutritional and pharmacological properties which will help studentsand researchers to get the overall information about its published nutritive and pharmacological properties for their further research.

Nutritional Composition

Most of the food components including macro- and micro-nutrients play important role as a nutraceutical, and provides potential health benefits (Bernal *et al*

2010). Dietary fiber, polyunsaturated fatty acids (PUFA), proteins, amino acids, minerals, vitamins and other bioactive compounds are considered as beneficial nutrient components (Andlauer and Fürst, 2002). Fresh ginger contains 80.9% moisture, 2.3% protein, 0.9% fat, 1.2% minerals, 2.4% fibre and

12.3% carbohydrates. The minerals present in ginger are iron, calcium and phosphorous. It also contains vitamins such as thiamine, riboflavin, niacin and vitamin C. The composition varies with the type, variety, agronomic conditions, curing methods, drying and storage conditions. [6]





Figure 1: Ginger plant and rhizome

Table 1:Nutritional composition of ginger (per 100g)

Constituent	Value	Constituent	Value
Moisture	15.02 ± 0.04	Ash (g)	$3.85 \pm 0.61 (4.53)$
Protein (g)	$5.087 \pm 0.09 (5.98)$	Calcium (mg)	88.4 ± 0.97 (104.02)
Fat (g)	$3.72 \pm 0.03 (4.37)$	Phosphorous (mg)	174±1.2 (204.75)
Insoluble fibre (%)	$23.5 \pm 0.06 (27.65)$	Iron (mg)	$8.0 \pm 0.2 (9.41)$
Soluble fibre (%)	$25.5 \pm 0.04 (30.0)$	Zinc (mg)	$0.92 \pm 0 \ (1.08)$
Carbohydrate (g)	38.35 ± 0.1	Copper (mg)	0.545±0.002 (0.641)
Vitamin C (mg)	$9.33 \pm 0.08 (10.97)$	Manganese (mg)	$9.13 \pm 001 (10.74)$
Total carotenoids (mg)	79 ± 0.2 (9296)	Chromium (µg)	$70 \pm 0 \ (83.37)$

All value in this table represent the mean \pm SD (n = 4). Figures in the parenthesis represent the dry weight values.

Chemistry

In the fresh ginger rhizome, the gingerols were identified as the major active components and gingerol [5-hydroxy-1-(4-hydroxy-3-methoxy phenyl) decan-3-one of the most abundant constituent in the gingerol series. The powdered rhizome

contains 3-6% fatty oil, 9% protein, 60-70% carbohydrates, 3-8% crude fiber, about 8% ash, 9-12% water and 2-3% volatile oil. The volatile oil consists of mainly mono and sesquiterpenes, camphene, beta-phellandrene, curcumene, cineole, geranyl acetate, terphineol, terpenes, borneol,

geraniol, limonene, linalool, alpha-zingiberene(30-70%), beta-sesquiphellandrene (15-20%), beta-bisabolene (10- 15%) and alpha-farmesene. In dried ginger powder, shogaol a dehydrated product of gingerol is a predominant pungent constituent.

Oleoresin, which is isolated by acetone and ethanol extraction, contains 4-7.5% of dried powder, pungent

substances namely gingerol, shogaol, zingerone and paradol. The oleoresin has also been found to contain zingiberol, the principal aroma contributing component as well as zingiberene, gingediol, diarylheptanoids, vitamins andphytosterols.^[7]

Figure 2: Chemical structure of active constituents of Zingiberofficinale.

Pharmacological Effects




Figure 3: Ginger and its constituents shows role in diseases prevention.

Anti-oxidant action

Ginger and some specific constituents have demonstrated antioxidant effects in several cell culture systems.[8-11] Furthermore, there are animal studies showing that ginger extracts and individual ginger constituents such as [6]-gingerol can protect several tissues and organs against damage due to a variety of oxidation-inducing stressors like ultra violet B (UVB) and COX-2 idiom a promising induced skin therapeutic agent against UVB disorders, has been studied both in-vitro & in-vivo. [12-^{19]} In rats, ginger extract also ameliorated acetic acid induced ulcerative colitis, likely due to its antioxidant and anti-inflammatory actions. [20] It also has a protective role to toxicity and lethality against some agent like carbon-tetra chloride, cisplatin etc. [21]

Anti- inflammatory

In Indian ayurvedic medicine, ginger is used as an anti-inflammatory herb. Ginger also shows a vital role in the suppression/inhibition in synthesis of pro-

inflammatory cytokines such as IL-1, TNF- α , and IL-8. $^{[22]}$

Anti tumor

Ginger also acts as antitumor via modulation of genetic pathways such as activation tumour suppressor gene, modulation of apoptosis and inhibition of VEGF.Study suggests that ginger extract may bring to bear its antitumor effects on colon cancer cells by suppressing its growth, striking the G0/G1- phase, reducing DNA synthesis and inducing apoptosis.^[23]

Effect on migraine

500-600mg of ginger powder administration at the onset of migraine for 3-4 days at interval of 4 hours, reported to provide relief from migraine attack^[24]

Neuro protective activity

The neuroprotective effect is partly attributable to an antagonistic action of ginger root extractson monosodium glutamate effect, so the monoamines content was increased. From these results, we can say

that the ginger extract has a neuroprotective role against monosodium glutamate toxicity effect [25]

Effect on Osteoarthritis

A highly purified and standardized ginger extract had a statistically significant effect on reducing symptoms of Osteoarthritis of the knee. This effect was moderate. There was a good safety profile, with mostly mild GI adverse events in the ginger extract group [26]

Hepato-protective effect

Earlier investigators based on experimental findings have shown that, ginger and its constituents play a significant role in hepato-protection. An important study on ginger showed its protective effect against the CCl4-induced hepatotoxicity. [27]

Anti- diabetic

Several animal studies indicate that ginger may be beneficial in lowering problematic blood glucose and lipid concentrations. Specific extracts of ginger lowered blood glucose, cholesterol, and triglyceride levels and increased high-density lipoprotein cholesterol concentrations.

Antimicrobial effects

Ginger has strong antibacterial and to some extent antifungal properties. In vitro studies have shown that active constituents of ginger inhibit multiplication of colon bacteria. These bacteria ferment undigested carbohydrates causing flatulence. This can be counteracted with ginger. It inhibits the growth of Escherichia coli, Proteus sp, Staphylococci, Streptococci and Salmonella.

Summary and Conclusion

The present study provides a strong base on the nutritional value of ginger with the presence of high amounts of dietary fiber, unsaturated fatty acids, essential amino acids and essential minerals with lower levels of heavy metal contaminants.

Based on the current body of scientific literature, ginger demonstrates some promising health benefits. It can be concluded that ginger shows an important effect in the suppression of NFkB, COX-2, and LOX, induction of apoptosis, activation of tumor suppressor gene and also modulates various biological activities. Ginger and their constituents create optimism towards the novel therapeutic strategy. Future research should focus on clinical trials to investigate its effectiveness and their exact role in modulation of molecular path-ways.

Reference

- 1. Kathi J. Kemper, MD, MPH. Ginger. Page 1. Longwood Herbal Task Force. Evans WC. London; Philadelphia: BaillièreTindall, 1989.
- 2. Bordia, A. Verma S.K. and Srivastava K.C. Prostaglandins.LeukotEssent Fatty Acids. 1997; 56: 379.
- 3. Srivastava K.C. Prostaglandins Leukot Med. 1984; 13: 227.
- **4.** Nutrition Today. Volume 45 Number 4 July/August. 2010.
- **5.** Ozgoli G1. Goli M.Moattar F. Comparison of effects of ginger, mefenamic acid and ibuprofen on pain in women with primary dysmenorrhea. J Altern Complement Med. 2009 Feb;15 (2):129-32.
- **6.** Gugnani H.C. Ezenwanze.Aeschbach R. Loliger J. Scott B. et al. Antioxidant actions of thymol, carvacrol, 6-gingerol, zingerone and hydroxytyrosol. Food Chem.Toxicol. 1994;32:31-36.16.
- **7.** Asnani V.Verma R. Antioxidative effect of rhizome of Zingiberofficinale on paraben induced lipid peroxidation: an in vitro study. Acta Pol Pharm. 2007;64:35-37.17.

- **8.** Chohan M. Forster-Wilkins G.Opara E. Determination of the antioxidant capacity of culinary herbs subjugated to various cooking and storage processes using the ABTS(+) radical cation assay. Plant Foods Hum Nutr. 2008;631:47-52.18.
- **9.** Tao Q. Xu Y. Lam R. et al. Diarylheptanoids and aminoterpenoid from the rhizome of Zingiberofficinale: antioxidant and cytoprotective properties. J Nat Prod. 2008;71:12-17.
- **10.** Kim J. Kim Y. Na K. Surh Y. Kim T. [6]-Gingerol prevents UVB-induced ROS production and COX-2 expression in vitro and in vivo. Free Radic Res. 2007;41:603-614.
- 11. Mallikarjuna K. Sahitya P. Sathyavelu K.Rajendra W. Ethanol toxicity: rehabilitation of hepatic defense system with dietary ginger. Fitoterapia. 2008;79:174-178.
- **12.** Ahmed R. Suke S. Seth V. Chakraborti A. Tripathi A. Banerjee B. Protective effects of dietary ginger (ZingiberofficinaleRosc.) on lindane-induced oxidative stress in rats. Phytother Res. 2008;22:902-906.
- **13.** Amin A.Hamza A. Effects of Roselle and ginger on cisplatin-induced reproductive toxicity in rats. Asian J Androl. 2006;8:607-612.
- **14.** Haksar A. Sharma A. Chawla R. et al. Zingiberofficinale exhibits behavioral radioprotection against radiation-induced CTA in gender-specific manner. PharmacolBiochemBehav. 2006;84:179-188.
- **15.** Ansari M. Bhandari U.Pillai K. EthanolicZingiberofficinale R. extract pretreatment alleviates isoproterenol-induced oxidative myocardial necrosis in rats. Indian J Exp Biol. 2006;44:892-897.
- **16.** Ajith T. Nivitha V.Usha S. Zingiber officinale Roscoe alone and in combination with alpha-tocopherol protect kidney against cisplatin-induced acute renal failure. Food ChemToxicol. 2007;45:921-927.
- **17.** Ajith T. Hema U.Aswathy M. Zingiber officinale Roscoe prevents acetaminophen-induced acute hepatotoxicity by enhancing hepatic antioxidant status. Food ChemToxicol. 2007;45:2267-2272.
- **18.** Ahmed R. Seth V. Pasha S.Banarjee B. Influence of dietary ginger (ZingiberofficinaleRosc.) on oxidative stress induced by malathion in rats. Food ChemToxicol. 2000;38:443-450.
- **19.** Zadeh J.B Kor N.H Physiological and pharmaceutical effects of Ginger (Zingiberofficinale Roscoe) as a valuable medicinal plant. Euro. J. Exp. Bio. 2014; 4(1):87-90
- **20.** El-Abhar H. Hammad L.Gawad H. Modulating effect of ginger extract on rats with ulcerative colitis. J Ethnopharmacol. 2008;118:367-372.
- **21.** Jagetia GC. Baliga MS. Venkatesh P. Ulloor JN. Influence of ginger rhizome (ZingiberofficinaleRosc.) on survival, glutathione and lipid peroxidation in mice after whole-body exposure to gamma radiation. Radiat Res. 2003; 160: 584–592.
- **22.** Tjendraputra E. Tran VH. Biu-Brennan D. Roufogalis BD. Duke CC. Effect of ginger constituents and synthetic analogues on cyclooxygenase-2 enzyme in intact cells. Bioorg. Chem. 2001; 29: 156-163.
- 23. Abdullah et al Ginger extract (Zingiberofficinale) triggers apoptosis and G0/G1 cells arrest in HCT 116 and HT 29 colon cancer cell lines. African Journal of Biochemistry Research. 2010; 4(4): 134-142.
- 24. Mustafa T and Srivastava KC, Ginger (Zingiberofficinale) in migraine headache. J Ethno. pharmacol. 1999; 29: 267.
- **25.** Waggas AM.Neuroprotective evaluation of extract of ginger (Zingiberofficinale) root in monosodium glutamate-induced toxicity in different brain areas male albino rats. Pak J Biol. Sci. 2009; 12(3): 201-212.

- **26.** Altman RD. Marcussen KC. Effects of a ginger extract on knee pain in patients with osteoarthritis. Arthritis Rehum. 2001; 44(11): 2531-2538.
- **27.** Patrick-Iwuanyanwu KC. Wegwu MO.Ayalogu EO. Prevention of CCl4- induced liver damage by ginger, garlic and vitamin E. Pak J Biol. Sci. 2007; 10: 617-62115.